



Content available at: <https://www.ipinnovative.com/open-access-journals>

Journal of Oral Medicine, Oral Surgery, Oral Pathology and Oral Radiology

Journal homepage: www.joooo.org



Original Research Article

Incidence of histopathological variants of oral squamous cell carcinoma: An institutional study

Saket Mehrotra^{1,*}

¹Dept. of Pathology, Sardar Patel Postgraduate Institute of Dental and Medical, Lucknow, Uttar Pradesh, India



ARTICLE INFO

Article history:

Received 26-10-2021

Accepted 25-11-2021

Available online 14-01-2022

Keywords:

Verrucous carcinoma (VC)

Adenoid/acantholytic/

pseudoglandular Squamous cell carcinoma (AdSCC)

Spindle cell/sarcomatoid carcinoma (SCSC)

Adenosquamous carcinoma (ASC)

Basaloid SCC (BSCC)

ABSTRACT

Background: Squamous cell carcinoma is the most important and the most common malignant mucosal neoplasm of the head and neck accounting for over 90% of all malignancies. Conventional oral Squamous cell carcinoma is frequently present in general cancerous conditions. It is bundled up with six different variants. Histomorphologically every variant shows a unique appearance. This raises an opportunity for the different diagnostic consideration with the precise management decision.

Methodology: All cases of OSCC reported at our institution Dentopath Pathologies Amravati in past two months were scrutinized. Representative sections containing the full thickness of the tumor were used for histopathological grading. The structure and identification of carcinomas were done microscopically by two expert dentopathologist.

Results: In the present study, we screened 100 slides of a conventional epithelial cell carcinoma. Amongst which 30 Slides showed the verrucous carcinoma. On 5 slides adenoid squamous cell carcinoma were observed. Incidence of Papillary squamous cell carcinoma and basaloid squamous cell carcinoma was only 1 out of 100 slides each. Whereas, the spindle cell/sarcomatoid carcinoma was observed on 2 slides. Adenosquamous carcinoma is the rarest variant and hence no incidence of this carcinoma were observed in our study.

Discussion: The behavior of the OSCC varies amongst due to the presence of different morphological type of tumor. A few studies on OSCC malignancy grading with different clinical parameters were made. In the present study different types of variants are seen according to their histopathological appearances.

Conclusion: Histopathological knowledge is very important for the precise diagnosis. Squamous cell carcinoma is the most common neoplasm of oral cavity. However, variants of the same show very less frequency. Hence, it became challenge for the appropriate diagnosis as many times a misdiagnosis affects the course of treatment of the patient

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Squamous cell carcinoma is the most important and the most common malignant mucosal neoplasm of the head and neck accounting for over 90% of all malignancies. Conventional oral Squamous cell carcinoma is frequently present in general cancerous conditions. It is bundled up with

six different variants namely verrucous carcinoma (VC), adenoid/ acantholytic/ pseudoglandular Squamous cell carcinoma (AdSCC), spindle cell/sarcomatoid carcinoma (SCSC), adenosquamous carcinoma (ASC), basaloid SCC (BSCC) and papillary Squamous cell carcinoma (PSCC).^{1,2} Histomorphologically every variant shows a unique appearance. Which raises an opportunity for the different diagnostic consideration with the precise management decision. It has been found that squamous cell carcinoma

* Corresponding author.

E-mail address: drsaket18@gmail.com (S. Mehrotra).

is usually arise due to oral cavities and common risk factors are over use of tobacco and alcohol consumption.² However, immunologic factors, age, gender, genetic factors and environmental conditions may also play a vital role. Viruses are also one of the developing factors for squamous cell carcinoma.

Verrucous carcinoma is the most common variant of squamous cell carcinoma seen in oral cavity, especially at the buccal mucosa or gingiva due to the excess use of tobacco, alcohol and poor oral hygiene.³ Clinically, VC of mouth composed of cauliflower like exophytic growth, which are thickened and club-shaped, and lined by well differentiated squamous epithelium.⁴ The microscopic examination of verrucous carcinoma appearance with abundant keratosis and parakeratosis rising with from a fold, acanthotic squamous epithelium which leads to appears like “church spires”.⁵ Parakeratotic crypting is the most common feature. Advancing margins of tumor are typically broad or bulbous rete pegs with pushing rather than an infiltrative appearance. Hybrid abrasion observe, when tumors present the dominant microscopic features of VC, but it contain small areas of tumor invasion.^{6,7} To diagnose VC, ample sectioning and good demonstrative sampling of the base of lesion are needed as evidence of invasion is required for definite diagnosis.⁸ The features of VC represent invasion with further acanthosis and broad based rete pegs prolonging deeply into the stromal tissue, while Verrucous hyperplasia should not spread more intensely than adjacent uninvolved epithelium. Also, Verruca Vulgaris has a prominent keratohyline granules and parakeratosis with sharply defined acanthotic rete ridges, not seen in VC. The biological behavior of VC is between non-neoplastic hyperplasia and conventional SCC.⁹ Therefore, it may be helpful to think of VC as an extremely well differentiated squamous cell carcinoma.

Adenosquamous cell carcinoma is a combination of squamous cell carcinomas and adenocarcinomas and is a high grade variant of SCC. These carcinomas spread throughout the upper digestive tract.¹⁰ The two carcinomas may be separate or intermixed, with areas of commingling or transition of the squamous cell carcinoma to adenocarcinoma. The undifferentiated areas between the two distinct carcinomas are often composed of clear cells. Both carcinomas may demonstrate frequent mitoses, necrosis and infiltration into the surrounding tissue with affiliated perineural invasion.^{11,12}

Basaloid Squamous Cell Carcinomas are variant of squamous cell carcinoma (SCC) of the head and neck. Tumor indicates variety of growth including solid, lobular, crebriform, cord, trabaculæ, nests and glands or cysts. Large nests can have central comedo form necrosis.¹³ The component basaloid is most diagnostic characteristic feature, incorporating small, closely opposed moderately pleomorphic cells with hyperchromatic nuclei and scant

cytoplasm into a lobular configuration with peripheral palisading, closely associated with or involving the surface mucosa.^{14,15} These basaloid regions are in direct continuity with areas of squamous differentiation including abrupt keratinization in the form of squamous pearls, individual cell keratinization, dysplasia or SCC.¹⁶

Papillary squamous cell carcinomas are the most uncommon and distinctive variants of squamous cell carcinomas. Papillary carcinomas by definition are without pre or co existing benign lesions.^{17,18} Papillary squamous cell carcinomas are uncommon but distinct variants of SCC of the upper aerodigestive tract mucosa, separable from verrucous SCC. The common most site for this infection is larynx followed by oro- and hypopharynx. Macroscopically, this carcinomas appearing as are polypoid, exophytic, bulky, papillary or fungiform tumours, soft to firm, arising from a broad base or from a narrow stalk. This carcinomas must demonstrate a dominant papillary architectural growth pattern with the evidence if malignancy. The papillary pattern comprises of multiple, thin, delicate filiform, finger-like papillary projections.¹⁹ The papillae contain a delicate fibrovascular core surrounded by the neoplastic epithelium. There are no specific or unique clinical or demographic parameters for this variant of SCC.²⁰

2. Materials and Methods

All cases of OSCC reported at our institution Dentopath Pathologies Amravati in past two months were scrutinized. Clinicopathological features of patients were compared. Clinical details were obtained from patients' records. Paraffin sections for the analysis of histopathological features were obtained from the archives of our department.

The parameters such as etiological factors, duration of risky habits, site and size of the tumor, morphological type, and histopathological grade of the tumor were examined. Representative sections containing the full thickness of the tumor were used for histopathological grading. The structure and identification of carcinomas were done microscopically by two expert dentopathologist.

3. Results

In the present study, we screened 100 slides of a conventional epithelial cell carcinoma. The large, bulbous downward proliferation was observed in 30 slides confirming verrucous carcinoma. On 5 slides adenoid squamous cell carcinoma were observed and was confirmed by the presence of lobular growth pattern with psuedoglandular alveolar areas. Incidence of Papillary squamous cell carcinoma was only 1 out of 100 slides which appeared as finger like projections with fibro vascular cores and bulbous growth with rounded projections. The basaloid squamous cell carcinoma on micro focusing was observed in 1 slide only. Microscopically it appeared as lobules, nests

or cribriform patterns with peripheral palisading and a thick basement membrane. Whereas the microcystic giant cells were observed on 2 slides emphasizing the presence of spindle cell/sarcomatoid carcinoma. (Figure 1) It has been seen that adenosquamous carcinoma is the rarest variant and hence no incidence of this carcinoma were observed in our study.




Fig. 1: Variant of conventional cell carcinoma

4. Discussion

The 90% of oral cancer accounts for conventional squamous cell carcinoma. Along with this carcinoma variants of OSCC frequently arise within the oral cavity. The behavior of the OSCC varies amongst due to the presence of different morphological type of tumor. A few studies on OSCC malignancy grading with different clinical parameters were made.^{2,5,16,21} In the present study different types of variants are seen according to their histopathological appearances. Joshua Franklyn et al.²² observed 1049 patients over 10 years of time period and supported the theory of 2-16% incidence rate of verrucous carcinoma. In the present study the carcinoma observed histopathologically and showed 30% of incidence, since the sample size of the present study is significantly lower. A study suggested that the lesion form due to these variant of OSCC are clinically difficult to differentiate and hence it is generally recommended to excise the lesion surgically. Basaloid squamous cell carcinoma is found to be in frequency of 1% of all squamous cell carcinoma which is in concordance with the study made by Thompson L who also concluded that the frequency of basaloid squamous cell carcinoma is >1%.⁵ The presence of spindle cell with elongated nuclei and scant eosinophilic cytoplasm at the periphery of the nest of the basoid cells is the microscopic appearance observed by Shinno Y et al.²³ in his case report which is similar to the microscopic observations found in our study. Yasunori T et al.²⁴ suggested that clinically, both papillary squamous cell carcinoma and verrucous carcinoma appear as an exophytic, papillary tumor, but the latter is white, warty and fungating with multiple filiform projections. Careful histopathological

investigation is needed to establish the correct diagnosis, which is essential for appropriate treatment their findings are concordance with our study.

Spindle cell carcinoma is a biphasic tumor with epitheloid and spindle shape neoplastic proliferation accounting for 3% of squamous cell carcinoma. This carcinoma is a rare tumor in the oral cavity accounting only 1% of all tumors of oral region. Findings of our study were similar to the cases reported to the Prakash N et al.²⁵ showing the similar cases of spindle cell carcinoma in the oral cavity involving alveolar ridge.

Adenoid squamous cell carcinoma is an uncommon and microscopically characterized by bundle of malignant epithelial cells leading to pseudoglandular appearance. Many studies emphasized on the unique histopathologic features as appeared in the present study that is the pseudoglandular appearance of carcinoma.²⁶

Adenosquamous cell carcinoma is histopathologically is a mixture of distinct components consisting of squamous cell carcinoma and adenosquamous cell carcinoma. Although, these two components are vary closing proximity. Our study findings, did not report any adenosquamous cell carcinoma concluding the rare nature of the variant.⁵

5. Conclusion

Histopathological knowledge is very important for the precise diagnosis. Squamous cell carcinoma is the most common neoplasm of oral cavity. However, variants of the same show very less frequency. Hence, it became challenge for the appropriate diagnosis as many times a misdiagnosis affects the course of treatment of the patient. For the dentopathologist to ensure the correct diagnosis multiple sectioning required. The conclusion made in the present study is limited to less sample size and in lesser population area. For the more firm conclusion a revise study with a different population area and bigger sample size is required.

6. Source of Funding

None.

7. Conflict of Interest

The authors declare no conflict of interest.

References

1. Thompson LD. Squamous cell carcinoma variants of the head and neck. *Curr Diagn Pathol*. 2003;9(6):384–96.
2. Pathak J, Swain N, Patel S, Poonja L. Histopathological variants of oral squamous cell carcinoma-institutional case reports. *J Oral Maxillofac Pathol*. 2014;18(1):143–5.
3. Spiro RH. Verrucous carcinoma, then and now. *Am J Surg*. 1998;176(5):393–7.
4. Kaugars GE, Abbey LM, Burns JC, Page DG, Svirsky JA. Oral Verrucous carcinoma. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 1999;87(2):268–77.

5. Thompson LDR. Squamous cell carcinoma variants of the head and neck. *Curr Diagn Pathol*. 2003;9(6):384–96.
6. Koch BB, Trask DK, Hoffman H. National survey of head and neck verrucous carcinoma: patterns of presentation, care, and outcome. *Cancer*. 2001;92(1):110–20.
7. Abramson AL, Brandsma J, Steinberg B, Winkler B. Verrucous carcinoma of the larynx. Possible human papillomavirus etiology. *Arch Otolaryngol*. 1985;111:709–15.
8. Thompson LD, Wenig BM, Heffner DK, Gnepp D. Exophytic and papillary squamous cell carcinomas of the larynx: A clinicopathologic series of 104 cases. *Otolaryngol Head Neck Surg*. 1999;120(5):718–24.
9. Batsakis JG, Hybels R, Crissman JD, Rice D. The pathology of head and neck tumours: verrucous carcinoma, Part 15. *Head Neck Surg*. 1982;5(1):29–38.
10. Gerughty RM, Hennigar GR, Brown F. Adenosquamous carcinoma of the nasal, oral and laryngeal cavities. A clinicopathologic survey of ten cases. *Cancer*. 1968;22(6):1140–55.
11. Keelawat S, Liu CZ, Roehm PC, Barnes L. Adenosquamous carcinoma of the upper aerodigestive tract: a clinicopathologic study of 12 cases and review of the literature. *Am J Otolaryngol*. 2002;23(3):160–8.
12. Damiani JM, Damiani KK, Hauck K, Hyams VJ. Mucoepidermoid adenosquamous carcinoma of the larynx and hypopharynx: a report of 21 cases and a review of the literature. *Otolaryngol Head Neck Surg*. 1981;89:235–43.
13. Stelow EB, Mills SE. Squamous cell carcinoma variants of the upper aerodigestive tract. *Am J Clin Pathol*. 2005;124(1):96–109.
14. Jayakrishnan A, Elmalah I, Hussain K. Basal cell adenocarcinoma in minor salivary glands. *Histopathology*. 2003;42(6):610–4.
15. Coletta RD, Cotrim P, Almeida OP. Basaloid squamous carcinoma of oral cavity: a histologic and immunohistochemical study. *Oral Oncol*. 2002;38(7):723–9.
16. Chovatiya NR, Modi TG, Chokshi VA, Shah S, Majeethia D. Oral Squamous Cell Carcinoma Variants - A Clinico-Pathologic Relevance. *J Dent Med Sci*. 2018;17(5):25–30.
17. Friedberg SA, Stagman R, Hass GM. Papillary lesions of the larynx in adults: a pathologic study. *Ann Otol Rhinol Laryngol*. 1971;80(5):683–92.
18. Ferrer MJ, Estelles E, Villanueva A. Papillary squamous cell carcinoma of the oropharynx. *Eur Arch Otorhinolaryngol*. 2003;260:444–5.
19. Ferlito A, Devaney KO, Rinaldo A. Papillary squamous cell carcinoma versus verrucous squamous cell carcinoma of the head and neck. *Ann Otol Rhinol Laryngol*. 1999;108(3):318–22.
20. Banks ER, Frierson HF, Mills SE, George E, Zarbo RJ, Swanson PE. Basaloid squamous cell carcinoma of the head and neck. A clinicopathologic and immunohistochemical study of 40 cases. *Am J Surg Pathol*. 1992;16(10):939–46.
21. Acharya S, Tayaar AS. Analysis of clinical and histopathological profiles of oral squamous cell carcinoma in young Indian adults: A retrospective study. *Journal of Dental Sciences*. 2012;7(3):224–30.
22. Franklyn J, Janakiraman R, Tirkey AJ, Thankachan C, Muthusami J. Oral Verrucous Carcinoma: Ten Year Experience from a Tertiary Care Hospital in India. *Indian J Med Paediatr Oncol*. 2017;38(04):452–5.
23. Shinnoy Y, Nagatsuka H, Siar CH, Tsujigiwa H, Tamamura R, Gunduz M. Basaloid squamous cell carcinoma of the tongue in a Japanese male patient: A case report. *Oral Oncol Extra*. 2005;41(4):65–9.
24. Nakanuma Y, Sato Y, Ojima H, Kanai Y, Aishima S, Yamamoto M, et al. Clinicopathological characterization of so-called "cholangiocarcinoma with intraductal papillary growth" with respect to "intraductal papillary neoplasm of bile duct (IPNB)". *Int J Clin Exp Pathol*. 2014;7(6):3112–22.
25. Prakash N, Kumar MS, Sharada P, Pradeep GL. Spindle cell carcinoma of the oral cavity: A case report of a rare entity and review of literature. *World J Dent*. 2010;(1):55–8.
26. Donthi D, Ramaswamy AS, Mahalingasetti PB. Acantholytic squamous cell carcinoma of the tongue: A diagnostic challenge. *Clin Cancer Investig J*. 2014;3(2):179–81.

Author biography

Saket Mehrotra, Reader and Head

Cite this article: Mehrotra S. Incidence of histopathological variants of oral squamous cell carcinoma: An institutional study. *J Oral Med, Oral Surg, Oral Pathol, Oral Radiol* 2021;7(4):226-229.