Case Report

Surgical excision of tooth embedded-osteoma of palate induced by traumatically intruded tooth – A case report

Ashwin V¹, Neha Sikka^{2,*}, Anbumani P³, Suman Bisla⁴, Narwade Pallavi Uddhav¹

¹Dept. of Oral and Maxillofacial Surgery, Pandit Bhagwat Dayal Sharma PGIDS, Rohtak, Haryana, India

²Dept. of Dental Materials, Pandit Bhagwat Dayal Sharma PGIDS, Rohtak, Haryana, India

³Dept. Oral and Maxillofacial Surgery, Chennai, Tamil Nadu, India

⁴Dept. of Oral Medicine and Radiology, Pandit Bhagwat Dayal Sharma PGIDS, Rohtak, Haryana, India

ARTICLE INFO

Article history:

Received 07-09-2020

Accepted 10-09-2020

Available online 03-10-2020

Keywords:

Hard palate

Peripheral osteoma

Trauma

Tntrusion

ABSTRACT

A 47-year old male patient reported with a chief complaint of hard swelling of palate region with associated difficulty in speech articulation and swallowing. It was diagnosed as osteoma of palate region, following a trauma with intrusion of tooth. Complete surgical excision was done and postoperative follow up revealed uneventful healing and restoration of articulation and tongue movements to near normalcy.

© 2020 Published by Innovative Publication. This is an open access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>)

1. Introduction

Osteomas are benign tumors of osteogenic origin. These tumors are slow in growth, round to oval in shape and commonly sessile, constituted of compact or cancellous bone. Various etiologic factors have been proposed for osteomas which include developmental, traumatic, infectious and neoplastic.^{1,2} They can be solitary or multiple. Gardner's syndrome is associated with multiple osteomas. Based on the origin, solitary osteomas can be classified as central, peripheral or extra-skeletal comprising soft tissue arising from the endosteum, periosteum or within a muscle respectively.^{3,4} The paranasal sinuses and the mandible are most commonly affected sites in the head and neck region.^{5,6} Palatal osteomas are relatively uncommon^{7,8} and are often asymptomatic. The osteomas in palatal region can cause functional impairment and minimal facial disfigurement. Surgical removal remains the gold standard treatment for osteomas with very low recurrence rate. In this article we present a case of pedunculated peripheral osteoma

of anterior palatal region in a 47 year old male, with its uniqueness lying on the traumatic intrusion of permanent tooth as etiologic factor. It was managed by surgical removal and primary closure.

2. Case Report

A 47 years male patient presented to our OPD with complaint of swelling in the region of palate since 25 years which was gradual in progress to attain the current size. The tumor was noticed by the patient following a trauma. Associated difficulty in chewing, restriction of tongue movements and speech articulation. There was no history of pain in the area of the swelling. Past medical history was non- contributory. On clinical examination, there was a large pedunculated swelling overlying the entire anterior part of hard palate. The swelling was approximately 3.5 * 4 cm, margins well defined and pedunculated with overlying mucosa appearing smooth and stretched. On palpation, there was no tenderness. The swelling was hard in consistency, non-mobile, non-fluctuant and there was clear demarcation between the swelling and the adjoining normal mucosa

* Corresponding author.

E-mail address: neha.nehasikka@gmail.com (N. Sikka).

(Figure 1). In addition to that there was missing central incisor on left side and fractured non vital lateral incisor and canine on the same side. With a provisional diagnosis of palatal exostoses and peripheral osteoma, radiographic investigations were undertaken. Occlusal view radiograph of upper arch revealed a well-defined radioopaque mass with a tooth like structure within it. Following which CECT was done, which showed a heterogenous, pedunculated, radio-opaque mass of 3* 4 * 3.5 cm, attached to the palate by a narrower stalk on the midline region. (Figure 2). Interestingly, a tooth like structure was present with proper crown and root morphology (Figure 2 B, D). This was correlating with the history of trauma and the absence of central incisor and fractured, intruded non-vital adjacent teeth. Based on clinical and radiological examination a provisional diagnosis of post-traumatic osteoma arising from the hard palate was made and excision biopsy was planned. Intraoperatively, a circumferential incision was given at the base to expose the mass (Figure 3 A). It was planned to preserve the firm and thick overlying mucosa as it will be helpful to cover the denuded region later. The osteotomy was started at the base using a tapered fissure bur to make the stalk thinner which was followed by separation from the hard palate using a curved osteotome and mallet (Figure 2 B) and the mass was excised in total (Figure 4). The remnants of stalk and the irregularities were smoothed in shaving motion using a trimming bur. The excess mucosa was reduced and primary closure was done with 3-0 vicryl following confirmation of intact palatal vault (Figure 3 C). The tumor was removed in toto and specimen showed a tooth engulfed within the tumor on the deeper surface. The postoperative period was uneventful and the wound healed completely (Figure 3 D). Histopathological examination showed that the lesion was composed of bony lamella with osteocytes and intervening fibrous tissue, fatty tissue and calcification. No cellular atypia was seen. Three years follow up showed no recurrence of the lesion.

Fig. 1: Preoperative photograph showing pedunculated tumor

Fig. 2: Preoperative radiograph: **A):** Sagittal section of CBCT; **B):** Occlusal view; **C):** 3D view; **D):** Note the displaced tooth within the tumor mass

Fig. 3: Intraoperative photograph: **A):** Exposure; **B):** Post excision; **C):** Closure; **D):** Follow-up photograph

3. Discussion

Osteomas are uncommon in palatal region and occur in a wide age distribution and without any gender predilection.³ According to study by Woldenberg et al, osteomas of

Fig. 4: Specimen. Note the discolouration representing the tooth engulfed within the tumor

maxillofacial region occurs in mean age group of 40.5 years.⁴ In this case the patient was a 47 year old male with history of trauma. Paranasal sinuses followed by mandible and maxillary region remain as the common site for peripheral osteomas occurring in the head and neck region. The hard palate is a rare site with a very few cases reported in literature. In this case, the trigger factor for onset of tumor was trauma. The intruded tooth into the anterior hard palate region following a trauma presented as if the tooth was buried within the bony mass. Radiographically, peripheral osteomas present as an oval to round, well circumscribed radio-opaque mass, and attached to the adjacent cortical bone with a broad base or pedunculated with a narrower stalk.⁸ Jaw bones exostoses as they do not grow beyond puberty, patients presenting with long standing and progressively growing bony masses should be examined for other lesions to rule out any associated syndromes. Asymptomatic osteomas or jaw bone exostoses are usually managed conservatively until they impede with function or esthetics. Complete surgical excision remains as gold standard treatment modality for symptomatic cases.⁹

4. Source of Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

5. Conflicts of Interest

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

6. Informed Consent

Written informed consent was obtained from a legally authorized representative for anonymized patient information to be published in this article.

References

1. Prabhuji MLV, Kishore HC, Sethna G, Moghe AG. Peripheral osteoma of the hard palate. *J Indian Soc Periodontol.* 2012;16(1):134–7.
2. Gundewar S, Kothari DS, Mokal NJ, Ghalme A. Osteomas of the craniofacial region: A case series and review of literature. *Indian J Plast Surg.* 2013;46(03):479–85.
3. Sayan NB, Üçok C, Karasu HA, Günhan Ö. Peripheral osteoma of the oral and maxillofacial region: A study of 35 new cases. *J Oral Maxillofac Surg.* 2002;60(11):1299–1301.
4. Woldenberg Y, Nash M, Bodner L. Peripheral osteoma of the maxillofacial region: diagnosis and management-a study of 14 cases. *Med Oral Patol Oral Cir Bucal.* 2005;10:139–42.
5. Larrea-Oyarbide N, Valmaseda-Castellón E, Berini-Aytés L, Gay-Escoda C. Osteomas of the craniofacial region. Review of 106 cases. *J Oral Pathol Med.* 2007;37(1):38–42.
6. Nah KS. Osteomas of the craniofacial region. *Imaging Sci Dent.* 2011;41(3):107–13.
7. Viswanatha B. Peripheral Osteoma of the Hard Palate. *Ear, Nose Throat J.* 2013;92(8):E31.
8. Ramoglu M, Yanik S, Bozdog Z, Aras MH. Peripheral osteoma of the hard palate: a rare case report and literature review. *J Istanbul Univ Fac Dent.* 2016;50(2):43–8.
9. Bountaniotis F, Melakopoulos I, Tzerbos F. Solitary Peripheral Osteoma of the Hard Palate: Case report and literature review. *Sultan Qaboos Univ Med J.* 2017;17(2):e234–7.

Author biography

Ashwin V Resident

Neha Sikka Assistant Professor

Anbumani P Consultant

Suman Bisla Resident

Narwade Pallavi Uddhav Resident

Cite this article: Ashwin V , Sikka N, Anbumani P , Bisla S, Uddhav NP. Surgical excision of tooth embedded-osteoma of palate induced by traumatically intruded tooth – A case report. *J Oral Med, Oral Surg, Oral Pathol, Oral Radiol* 2020;6(3):163-165.